

## C-SECTION SSI BUNDLE

<b>Pre Operative</b>
<b>Clinic</b>
<input type="checkbox"/> Chlorhexidine shower the night before and morning of surgery
<b>Pre Op Area Day of Surgery</b>
<input type="checkbox"/> Pre-operative warming (maintain temp >36.0)
<input type="checkbox"/> Full body Chlorhexidine (SAGE) wipes if showers not done
<input type="checkbox"/> Clip hair to 1 inch below surgical incision
<input type="checkbox"/> Check blood sugars for ALL diabetic patients (Goal 70 to 120)
<b>Intra Operative</b>
<input type="checkbox"/> Limit traffic into and out of OR, ensure appropriate OR attire
<input type="checkbox"/> ChlorPrep skin prep
<input type="checkbox"/> Iodine vaginal prep
<input type="checkbox"/> Antibiotic dosing within 1 hour of incision: Cefazolin: <120kg – 2gm IV, >120kg – 3gm IV <b>Add</b> Azithromycin (If active labor or ROM): 500mg IV over 1 hour Anaphylactic PCN Allergy: Clinda 600mg IV & Gent 5mg/kg IV
<input type="checkbox"/> If EBL or QBL > 1500 cc's, re-dose abx
<input type="checkbox"/> Placenta removal by traction on cord
<input type="checkbox"/> Change full set of gloves prior to starting fascial closure
<input type="checkbox"/> Maintain normothermia – OR temperature 72 degrees
<input type="checkbox"/> Use sutures for skin closure, close subcutaneous tissue if >2 cm
<input type="checkbox"/> Label dressing with date & time of placement
<b>Post Operative</b>
<input type="checkbox"/> Wound care education in PACU - patient and family
<input type="checkbox"/> Leave dressing on for 24 hours

Created July 2017

Updated August 2019

