





Inpatient Donning Checklist (N95)			Visuals
Read each step prior to beginning			
Step Name	✓		
Supplies		<b>Gather all supplies needed for donning and doffing</b> <ul style="list-style-type: none"> <li>In room supplies (initial room set up): hospital approved disinfectant wipes, hand sanitizer, trash can</li> <li>Outside room supplies: hospital approved disinfectant wipes, hand sanitizer, foot pedal trash can</li> </ul>	
Pre-Donning Activities		Check that the Donner:	
		<ul style="list-style-type: none"> <li>Changed into green hospital issued scrubs if indicated</li> </ul>	
		<ul style="list-style-type: none"> <li>Removed all personal jewelry, like earrings or watches that may interfere with doffing</li> </ul>	
		<ul style="list-style-type: none"> <li>Used the restroom</li> </ul>	
		<ul style="list-style-type: none"> <li>Hair is pulled back/controlled</li> <li>Hydrated</li> </ul>	
		<b>Donner:</b>	
Hand Sanitize		<ul style="list-style-type: none"> <li>Hand sanitize with Alcohol-Based Hand Sanitizer (ABHS)</li> </ul>	
Donning Standard Isolation Gown		<ul style="list-style-type: none"> <li>Don the over the head isolation gown in your size</li> </ul>	
		<ul style="list-style-type: none"> <li>Thumb should be inserted into the designated thumb loops</li> </ul>	
		<ul style="list-style-type: none"> <li>The front and back of gown should be pulled down for complete coverage</li> <li>The gown should fully cover torso from neck to knees, arms to end of wrists, and wrap completely around the back.</li> </ul> <p>For donning and doffing instructions please see the IPCD webpage:  <a href="https://hospitals.health.unm.edu/ipcd/wp-content/uploads/sites/4/2018/11/don.doff_.pdf">https://hospitals.health.unm.edu/ipcd/wp-content/uploads/sites/4/2018/11/don.doff_.pdf</a> </p>	
		<ul style="list-style-type: none"> <li>The side ties should be tied snug, but not too tight</li> </ul>	
Don Gloves		<ul style="list-style-type: none"> <li>Don gloves, ensure gloves extend over cuff of gown.</li> </ul>	
Donning N95 Mask		<ul style="list-style-type: none"> <li>Obtain the appropriate N95 respirator (indicated by your fit test)</li> </ul>	
		<ul style="list-style-type: none"> <li>Don N95 Mask:</li> <li>If donning a new N95 follow CDC Donning Guidance below.</li> <li><b>If following the Extended use and Limited Re-use PPE Guidance, please follow the process for N95 mask retrieval and then follow the steps for donning listed below.</b></li> </ul> <p><a href="https://www.cdc.gov/vhf/ebola/hcp/ppe-training/n95respirator_coveralls/donning_09.html">https://www.cdc.gov/vhf/ebola/hcp/ppe-training/n95respirator_coveralls/donning_09.html</a></p>	
		<b>Limited Re-Use Mask Retrieval</b> <ul style="list-style-type: none"> <li>Remove N95 mask from storage area and visually inspect for distortion. If it creases or bent, do not re-use</li> <li>Don the N95 Respirator following the CDC Donning steps below</li> </ul>	

	<b><u>CDC Donning Guidance:</u></b> <ul style="list-style-type: none"> <li>Put on the N95 respirator. Hold the respirator in the palm of your hand with the straps facing floor</li> <li>Place the N95 respirator on your face covering your nose and mouth</li> <li>Pull the bottom strap up and over top of your head, and put it behind your head below your ears</li> <li>Take the upper strap and put it behind your head towards the crown of your head</li> <li>Mold the nose piece of the respirator over the bridge of your nose to obtain a tight seal</li> </ul>	
<b>Negative/Positive Seal check</b>	<ul style="list-style-type: none"> <li>Once donned, perform a negative/positive seal check:               <ul style="list-style-type: none"> <li>No air should be felt around the perimeter while blowing out. If you feel air coming out it is not a tight seal.</li> <li>When taking a small breath in, the mask should pucker in slightly. If it does not, it is not re-usable</li> <li>When breathing out you should feel the respirator expand slightly. If it does not, it is no re-usable</li> <li>If not a tight seal, the respirator cannot be reused and should be placed in ppe collection bin for bioquell</li> </ul> </li> <li>Ensure the mask is breathable, if unable to breathe in the mask, the respirator cannot be re used</li> </ul>	
<b>Clean gloves</b>	<ul style="list-style-type: none"> <li>Sanitize gloves with Alcohol-Based Hand Sanitizer (ABHS)</li> </ul>	
<b>Donning a reusable face shield <u>OR</u> TIDI Shield/Goggles + procedure mask</b>	<p><b>If assembling the new, reusable face shield:</b></p> <ul style="list-style-type: none"> <li>Remove blue protective peeling from both front and back of shield</li> <li>Attach to the head band using the small holes on either side</li> </ul> <p><b>Donning assembled face shield:</b></p> <ul style="list-style-type: none"> <li>Don face shield so it sits level on your head</li> </ul> <p><b>Donning TIDI Shield</b></p> <ul style="list-style-type: none"> <li>Don TIDI Shield or Goggles so they sit level on your head</li> <li>Don procedure mask <i>over</i> N95</li> </ul>	
<b>Bouffant</b>	<p>Bouffant Cap:</p> <ul style="list-style-type: none"> <li>Apply Bouffant Cap (two may be necessary for longer hair)</li> <li>Make sure ears and forehead are completely covered</li> </ul>	
<b>Clean gloves</b>	<ul style="list-style-type: none"> <li>Sanitize gloves with Alcohol-Based Hand Sanitizer (ABHS)</li> </ul>	
<b>PPE check</b>	<ul style="list-style-type: none"> <li>Complete final check to ensure PPE correctly applied (360° view)</li> </ul>	
<b>Patient Room Reminders</b>	<ul style="list-style-type: none"> <li>Remember to move slowly and avoid reflexive actions such as touching your face</li> </ul>	
	<ul style="list-style-type: none"> <li>Use hand sanitizer when you doff your gloves, before donning on new gloves.</li> </ul>	
<b>Room Entry</b>	<ul style="list-style-type: none"> <li>You are ready to either enter the patient's room or assist with another HCWs doffing procedure</li> </ul>	